

ADVANCES IN INSTRUMENTATION AND TECHNIQUES IN STEM CELLS RESEARCH: A COMPREHENSIVE LITERATURE REVIEW

Sunil Kumar¹, Arti Thakur²

¹Origin Life Chandigarh, Department of Clinical Embryology & Reproductive Genetics, Rayat Bahara University, Mohali, Punjab, India

²Associate Professor cum HoD, Faculty of Allied and Healthcare Sciences, Desh Bhagat University, Gobindgarh, Punjab, India

Journal	International Research & Advancement in Microbiology and Biotechnology https://www.samvaktijournals.com/iramb Volume 1 Issue 1 (2025); Page No : 63 - 92
Discipline	Microbial Pathogenesis
Date Received	: July 31, 2025
ID	: iramb.2024.13
DOI No.	: 10.46402/iramb.2025.13
Publication Date	: December 30, 2025
Paper Type	: Review Paper
DOI Url	: https://dx.doi.org/10.46402/iramb.2025.13

ABSTRACT

Stem-cell research is an essential basis of modern biomedical research, given the unique and remarkable properties of stem cells. This literature review assesses the currently available knowledge of stem cells, while also acknowledging the continuously changing landscape of advanced technologies and instrumentation techniques in the discovery of stem cell structure and function. It also provides a thorough appraisal of the complexity of stem cell culture, which includes protocols and culturing conditions that maintain pluripotency and cell viability.

The review provides a snapshot of modern genetic approaches, neural CRISPR/Cas9, used with conventional transfection and transduction approaches, to make specific cell phenotypes. Advances in imaging, single cell approaches, and live cell tracking also provide a close-up look at stem cell behaviours whereas biomaterials and scaffolds supports its growth and engineering. Clinical applications of stem cells are also noted through current trials related to disease modelling and regenerative therapies. Importantly, ethical, regulatory and technical issues including reproducibility and scaling issues were carefully considered. This narrative provides information and direction to both experienced and inexperienced researchers, embryologists,

clinicians and trainees, by highlighting progress and acknowledging issues within the rapidly-evolving world of stem cell research.

Keywords: Stem cells; CRISPR/Cas9; Personalized Medicine; Regenerative Therapies; Single-Cell Analysis; Biomaterials.

INTRODUCTION

Background and Importance

Their capacity for sustained proliferation and their potential to develop into specialized somatic cells have positioned them as a major area of focus in contemporary biomedical research. They are distinguished by their remarkable capacity to give rise to multiple specialized cell lineages, making them indispensable to advances in basic science as well as translational biomedical research. The main types

1. “Embryonic stem cells” (ESCs)
2. “Adult stem cells (ASCs)”, and
3. “Induced pluripotent stem cells” (iPSCs)
4. Embryonic stem cells

arise from the inner cell mass of the blastocyst stage of the embryo and are pluripotent. As seen in *Figure 1* Embryonic stem cells arise from the inner cell mass of the blastocyst stage of the embryo and can become almost all cells in the body. However, pluripotent ESCs raise ethical concerns and possess the capacity to form tetrameric formations when used in therapeutic functions. Distributed among various tissues such as bone marrow, liver, and brain, adult stem cells exhibit multipotency, allowing them to contribute significantly to the ongoing maintenance and regeneration of the tissues in which they reside (2). Today, adult stem cells have fewer ethical concerns and are used more frequently in therapeutics(3). “Induced pluripotent stem cells (iPSCs) are derived by converting mature somatic cells into a pluripotent state through the ectopic activation of a defined combination of transcription factors, and combine the benefit of ESCs type pluripotency with the reduced ethical issues of ASCs (4). iPSCs carry the potential for personalized clinical applications, in addition to disease modelling and providing a powerful platform for Regeneration-focused therapeutic strategies. The importance of stem cells in biomedical approaches aimed at tissue repair and renewal can hardly be exaggerated. They offer a possible solution to degenerative diseases and trauma which fall short of current conventional medical therapies. For example, Ongoing research endeavors directed toward the development of stem cell-based therapeutic approaches for Parkinson and diabetes disease (5). In addition, stem cell-based models provide powerful tools for dissecting disease pathophysiology and

supporting the development and evaluation of novel therapeutic agents. Advances in stem cell-derived organoid technology have opened new avenues for studying organogenesis, disease dynamics, and treatment responsiveness within highly controlled model systems. (6).




Figure 1 : Illustration of Stem Cell Types

SCOPE AND OBJECTIVES

This article will present an in-depth description of the instrumentation and important procedures involved with stem cell science and applications. We chose to concentrate our efforts on summarizing and reviewing the instrumental developments that have ushered the science of stem cells into the current era, with an emphasis on more accurate study of properties and possibilities of the science. This article considers the gamut of methods and techniques in stem cell research from the basic concepts and protocols of stem cell culture to advanced methods in genetic engineering and microscopy. The article discusses major instruments (e.g., microscopy, flow cytometry, mass spectrometry, and next-generation sequencing (NGS)) needed to characterize the stem cell populations, deciphering the molecular signatures, modulating their functions *in vitro*, and tracking cellular behavior *in vivo*. Notably, this review includes novel developments in stem cell research, including CRISPR/Cas9 genome editing, high-content screening as a platform for drug discovery, and bioprinting in tissue engineering. Every topic discusses their ramifications, advantages, and their limitations, as well as the basics of the method itself. With this review, we provide an extensive outline of the tools and techniques of the field, as we hope to add to the developing knowledge base of researchers to enhance their ability

to leverage the use of these technologies for their research. It showcases emerging trends and current advancements, whilst suggesting ways these discoveries may impact the future of stem cell research. The primary purpose is to marry emerging technology with novel applications and to familiarize readers with the state of the art and future directions in stem cell science's instrumentation and technique.

LITERATURE REVIEW

Stem Cell Biology: Basic Concepts

Properties and Characteristics

Stem cells demonstrate a unique capacity to sustain continuous self-renewing activity and to differentiate across diverse specialized cellular lineages, making them highly valuable in both fundamental biological investigation and clinical practice. In this self-renewal process, stem cells undergo division to produce identical progeny that retain an undifferentiated state, thereby ensuring a sustained pool of stem cells. This self-renewal process includes "Wnt, Notch, and Hedgehog" signalling pathways, that act in a complex manner to ensure its characteristics are kept in a distinct balance (6). The differentiation process is where the stem cell is progressively converted through a series of regulated changes into a specialized cell type. This process is influenced by intrinsic and/or extrinsic factors; intrinsic factors including patterns of gene activity and epigenetic regulation, and extrinsic factors including cues derived from the surrounding microenvironment. During cellular differentiation, the stem cells progressively lose their self-renewal capability, while simultaneously gaining the ability to carry out specialized functions.

Stem Cell Niches and Microenvironments

Stem cell behavior is largely affected by the niches in which they exist. These niches contain the specialized microenvironments with physical and biochemical signals required to maintain stem cell properties and regulate their fate decisions, which can be seen in *Table 1*. Whereas the cellular microenvironment associated with a stem cell niche is shaped through the interactions between distinct cell types present within the niche, which can provide factors through direct contact with the cells as well as para-crine signalling. Moreover, extracellular matrix (ECM) elements including collagen fibres, lamina, as well as fibronectin provide mechanical support and are devising roles in signalling via integrin receptors and other ECM-binding proteins. Hypoxia, or a low oxygen condition, is another important aspect in some stem cell niches which regulates stem cell function through hypoxia-inducible factors (HIFs) and supports a self-renewal state while inhibiting differentiation (13). Cancer stem cells

(CSCs) are thought to exploit normal stem cell niches enhancing the microenvironment to support the growth of tumors and resistance to therapy (14). Therefore, knowledge of the complexities of stem cell niches and regulatory elements is essential for developing strategies to manipulate stem cell behavior for therapeutic manipulation as well as to devise interventions to regress CSCs in cancer therapy.

Section	Subsection	Key Points
Stem Cell Properties and Characteristics	Self-renewal	<ul style="list-style-type: none"> - Stem cells divide to produce more stem cells. - Maintained by signalling pathways.
	Differentiation	<ul style="list-style-type: none"> - Process of stem cells becoming specialized cells. - Influenced by intrinsic and extrinsic factors. - Hematopoietic differentiation.
	Types of Stem Cells	<ul style="list-style-type: none"> - ESCs: Pluripotent, derived from embryos. - ASCs: Multipotent, found in adult tissues. - iPSCs: Reprogrammed from somatic cells.
Stem Cell Niches and Microenvironments	Cellular Microenvironment	<ul style="list-style-type: none"> - Interactions with stromal, immune, and endothelial cells. - Example: Bone marrow niche and HSC maintenance.
	Extracellular Matrix (ECM)	<ul style="list-style-type: none"> - Provides structural support and active signalling. - ECM components: Collagen, laminin, fibronectin.
	Molecular Microenvironment	<ul style="list-style-type: none"> - Gradients of soluble factors: growth factors, cytokines, chemokines. - Example: Wnt signalling in stem cell regulation.
	Hypoxia	<ul style="list-style-type: none"> - Low oxygen conditions influence stem cell function. - Hypoxia-inducible factors (HIFs) promote self-renewal.
	Disruptions and Disease	<ul style="list-style-type: none"> - Niche disruptions can lead to diseases. - Cancer stem cells exploit niches for tumour growth.

Table 1 : Summary of Key Concepts in Stem Cell Properties, Characteristics, and Niches

Key Instrumentation and techniques in Stem Cell Research (illustrated in Figure 2)

Microscopy Techniques

Microscopy is a cornerstone within stem cell biology as it enables the visualization of cellular structures, the monitoring of dynamic processes, and the analysis of molecular interactions. Fluorescence microscopy is a widely adopted method of choice because of its ability to visualize biomolecules and cells by tagging them with a fluorescent dye or fluorescent protein to identify a specific protein of interest, organelle, and other cellular components. This method is useful for the examination of morphology, cell

fate specification (differentiation) and interactions with the stem cell niche (15). Confocal microscopy adds to fluorescence microscopy through the use of a pinhole aperture that eliminates out-of-focus light, producing sharper and more detailed images. This method is highly advantageous in producing the greatly sought-after high-resolution, three-dimensional reconstructions that stem cell researchers expect in studies of stem cells within their niches (16). Live-cell imaging allows researchers to observe stem cells as they behave in real-time, and visualize the STEM cell division, migration and differentiation under various spatially- and temporally-controlled conditions.

This step forward has transformed the field by enabling boxed visualization of subcellular structures and molecular complexes in stem cells and has led the way to better understanding the operating mechanisms in stem cell function and fate (17).

Flow Cytometry

To examine the physical and chemical properties of cell this is an analytical approach. With respect to stem cells, this technique serves as a highly effective tool for characterizing cellular populations based on surface and intracellular marker expression. Flow cytometry can quickly and quantitatively apply size, granularity, and fluorescence intensity to understand various cellular characteristics (18). Fluorescence-Activated Cell Sorting (FACS) can be used to separate required cell populations from the combination of multiple population based on fluorescence labelling. FACS is very useful for purifying stem cells and differentiating cells for further experiments, while allowing detailed investigation of defined cell types or developmental stages (19).

Mass Spectrometry

Mass spectrometry (MS) has become a significant tool in stem cell research for both proteomics and metabolomics. With MS, we can identify and quantify proteins, metabolites, and other biomolecules, and provide global profiles of stem cell states and states of response to varied stimuli. In proteomics, MS exposes differential protein composition as well as post-translational modifications and interactions amongst stem cells, and thus informs on cellular processes at the molecular level (20). Metabolomics uses MS to analyze the metabolic profiles of stem cells, providing insights into their metabolic states and pathways during differentiation and self-renewal processes (21).

Next-Generation Sequencing (NGS)

This technology has fundamentally transformed genomic and transcriptomic analysis in stem cell research.. NGS enables rapid and high-throughput sequencing of RNA

and DNA, and has heightened resolution for examining the genetic and transcriptomic scopes pertinent to stem cell research. This technique allows for the identification of gene expression patterns, regulatory elements and genetic mutations that drive stem cell differentiation and behaviour (22). In addition, researchers can study the heterogeneity in stem cell populations by using NGS to carry out single-cell RNA sequencing, allowing the characterization of subpopulations that have unique functional attributes and molecular signatures (23).

High-Content Screening

It integrates automated imaging with sophisticated data analysis to enable large-scale screening of stem cells. HCS is used to analyze large populations of cells across differentiated conditions to obtain quantitative data on cellular phenotypes, morphology, and marker expression patterns. This technology is primarily applicable in drug development to identify compounds that improve or inhibit patterns of stem cell proliferation, differentiation, or survival (24). Due to its automated and scalable nature of HCS, this approach has become a very important tool for many studies, which include new discovery of therapeutic target and optimization of stem cell-based therapeutics.




Figure 2 : Key Instrumentation in Stem Cell Research

Section	Subsection	Key Points	Reference
Microscopy Techniques	Fluorescence Microscopy	<ul style="list-style-type: none"> - Visualization of specific proteins and organelles. - Utilizes fluorescent dyes/proteins. 	(15)

Section	Subsection	Key Points	Reference
	Confocal Microscopy	<ul style="list-style-type: none"> - Eliminates out-of-focus light for clearer images. - Enables 3D reconstruction of stem cells. 	(16)
	Live-Cell Imaging	<ul style="list-style-type: none"> - Live monitoring of stem cells. - Monitors cell division, migration, differentiation. 	(25)
	Super-Resolution Microscopy	<ul style="list-style-type: none"> - Overcomes the diffraction limit of conventional light microscopy. - Detailed visualization at nan scale. 	(17)
Flow Cytometry	Principles and Applications	<ul style="list-style-type: none"> - Measures cell characteristics via light beam. - Analyses cell size, granularity, fluorescence. 	(14)
	Sorting Techniques (FACS)	<ul style="list-style-type: none"> - Isolates specific cell populations based on fluorescence. - Used for purifying stem cells. 	(19)
Mass Spectrometry	Proteomics	<ul style="list-style-type: none"> - Identifies and quantifies proteins. - Reveals protein composition and modifications. 	(20)
	Metabolomics	<ul style="list-style-type: none"> - Analyses metabolic profiles. - Offers insights into metabolic states during differentiation. 	(21)
Next-Generation Sequencing (NGS)	Genomic Analysis	<ul style="list-style-type: none"> - High-throughput DNA sequencing. - Identifies gene expression patterns, regulatory elements. 	(22)
	Transcriptomic Analysis	<ul style="list-style-type: none"> - RNA sequencing for detailed transcriptomic landscapes. - Uncovers subpopulations within stem cells. 	(23)
High-Content Screening	Automated Imaging and Analysis	<ul style="list-style-type: none"> - High-throughput screening of stem cells. - Analyses cellular phenotypes, morphology, marker expression. 	(24)

Table 2 : Key Instrumentation

Stem Cell Culture & Maintenance

Culture Conditions and Media

The successful culture of stem cells relies on careful optimization of the culture conditions and media as shown in *Table 3*. The basic components of stem cell culture media are complementation of basal media with nutrients such as vitamins, amino acids, sodium, glucose, and growth factors and cytokines that promote cell

proliferation and pluripotency. One of the most important supplements is serum, preferably fetal bovine serum (FBS), with a complex mix of growth factors, hormones, and adhesion factors. However, serum use can introduce variability and potential contamination risk, resulting in serum-free and chemically defined media (26). Chemically defined media create a controlled environment, which reduces variability and increases reproducibility in stem cell culture (27). Feeder layers usually consist of mitotically inactive murine embryonic fibroblasts (MEFs) which helps in the sustained proliferation of human embryonic stem cell populations (hESCs). Feeder layers also offer a supportive extracellular matrix and essential growth factor secretion, but the use of feeder layers from animals gave rise to potential cross species animal contamination and immunogenicity and resulted in the development of systems that do not need a feeder layer. Feeder-free culture systems use substrates such as Matrigel, which is a reconstituted basement membrane matrix, or synthetic matrices such as vitronectin and laminin and defined media to maintain pluripotency and promote stem-cell self-renewal (29). The developments have enabled the formation of xeno-free culture conditions that are fundamental for the translational clinical applications of therapies from stem-cell derivatives (27).

Cryopreservation and Thawing

Cryopreservation is a crucial approach for the long-term storage of stem cells, enabling researchers to preserve valuable cell lines and retain genetic and phenotypic stability over time (depicted in *Table 3*). Cryopreservation refers to the procedure of cooling the cells to sub-zero degrees, usually in liquid nitrogen (-196°C), while also introducing cryoprotective agents (CPAs) such as dimethyl sulfoxide (DMSO), and glycerol, which helps to protect cellular structures from the formation of ice crystals and prevent some degree of cellular damage during the freezing and thawing process (28). The most effective cryopreservation protocol utilizes a controlled-rate freezing procedure, which cools the cells slowly to help alleviate thermal shock and ice crystal formation (30). Thawing is also equally important and must also be performed quickly, to reduce the amount of time the cells are at intermediate temperatures and avoid the toxicity of the CPAs. Following thawing the cryovials are quickly placed into a 37°C water bath until thawed and subsequently, the CPAs are diluted to minimize cytotoxicity (31). The recovery and viability of post-thaw cells is greatly influenced by the thawing technique. Therefore, protocols must be meticulously optimized to promote viability and recovery, as well as for maintenance of cell functionality.

Passaging and Expansion

Passaging (also known as subculturing) is the move of cells from a dense culture vessel to a fresh culture vessel to allow more space for continued growth (see *Table*

3). Performing this procedure is critical to ensure that stem cells can remain viable and pluripotent for extended periods. Cells can be detached from the culture surface by enzymatic dissociation methods that utilize the enzymes trypsin or collagenase. However, these enzymes may damage cell surface proteins (marker proteins) which are critical to stem cell maintenance, which is why less damaging non-enzymatic dissociation solutions are used (32). Mechanical dissociation methods (e.g., scraping or pipetting) are an alternative but are generally less efficient and may induce cell stress than enzymatic methods. A "good" passaging protocol provides a balance between effectiveness of detaching cells from culture and minimizing affect on cell health. If more stem cells are to be expanded, particularly, if stem cells are to be produced at scale for potential therapeutic applications, suspension culture and bioreactor systems will be required. These systems must offer a controlled environment for expanding stem cells, while maintaining their undifferentiated state and functionality (33). Use of bioreactors has numerous advantages, namely with the ability to tightly control the culture environment, reduce human manipulation, and mechanise scaling up stem cell manufacture. Modulators of signalling pathways further support the maintenance of cell viability and pluripotency during passaging and expansion. ROCK inhibitors are one example of signalling pathway modulators which will enhance cell survival during single-cell dissociation of human pluripotent stem cells (hPSCs) since these cells are very sensitive to apoptosis when cultured as single cells (32). Advancements in techniques and technologies in cell culture is critical for reliable and reproducible expansion of stem cells so that they can be used for research and clinical application.

Section	Subsection	Key Points	References
Culture Conditions and Media	Essential Components and Optimization	<ul style="list-style-type: none">-Basal media supplemented with amino acids, vitamins, glucose, and salts.-Use of growth factors and cytokines to sustain proliferation and pluripotency.-Development of serum-free and chemically defined media for controlled environments.	(26, 27)
	Feeder Layers and Feeder-Free Systems	<ul style="list-style-type: none">-Feeder layers (e.g., MEFs) provide support and essential growth factors.	(29) (27)

Section	Subsection	Key Points	References
		<ul style="list-style-type: none"> -Feeder-free systems use substrates like Matrigel and synthetic matrices (e.g., vitronectin, laminin). -Xeno-free culture conditions for clinical applications. 	
Cryopreservation and Thawing	Methods and Protocols for Long-term Storage	<ul style="list-style-type: none"> -Use of cryoprotective agents (CPAs) like DMSO and glycerol to prevent ice crystal formation. -Controlled-rate freezing for gradual cooling. -Rapid thawing in 37°C water bath followed by CPA dilution. 	(28, 31)
Passaging and Expansion	Techniques for Maintaining Cell Viability and Pluripotency	<ul style="list-style-type: none"> -Enzymatic dissociation with trypsin or collagenase; use of non-enzymatic solutions for gentler cell detachment. -Mechanical dissociation methods (scraping, pipetting). -Use of bioreactors for large-scale production and controlled environments. -ROCK inhibitors to enhance cell survival during single-cell dissociation. 	(32, 33)

Table 3 : Techniques for Stem Cell Culture and Maintenance

Differentiation Techniques

Protocols for Directed Differentiation

Directed differentiation procedures are important for differentiating pluripotent stem cells into discrete cell lineages that serve a prominent role in both basic research and in therapeutic applications. Directed differentiation procedures often recapitulate many of the natural developmental cues that occur during early embryonic development. In most approaches, researchers use a combination of signaling cues, bioactive compounds, and/or extracellular matrix components to direct stem cells through a series of developmental events. This sequence design is representative of both

mesodermal induction and cardiac specification, similar to the process that occurs during these events *in vivo*. The application of regulatory growth cues and signal transduction pathways is an important component of the controlled differentiation strategy. Various regulatory growth mediators such as “fibroblast growth factor” (FGF), “epidermal growth factor” (EGF), and “transforming growth factor-beta” (TGF- β) are the primary categories of signalling molecules employed to promote stem cell differentiation into distinct cell types. As an example, differentiating neural progenitors from hPSCs requires the addition of FGF and EGF to regulate the proliferation and survival of neural precursors (34). Cell fate determination is influenced by various signaling pathways including Wnt, Notch, and Hedgehog, as precise regulation of these pathways can enhance differentiation efficiency and fidelity of differentiation protocols (35).

Organoid and 3D Culture Systems

The creation of organoid-based and three-dimensional culture platforms presents a major breakthrough in stem cell research that gives rise to more physiologically relevant models than conventional two-dimensional cultures. Organoids are three-dimensional, self-organizing constructions that are produced from stem cells and capture essential features of their respective organs. They provide an invaluable platform for investigating organogenesis, disease modelling, and drug screening with close resemblance to *in vivo* conditions (36). For instance, human embryonic stem cell-derived cerebral organoids were employed to simulate brain development and neurodevelopmental diseases and to elucidate the pathogenesis of diseases (36). These systems employ scaffolds composed of biocompatible materials like hydrogels to enable the growth and organization of stem cells into tissues with a high degree of complexity. Hydrogels may be engineered to mimic the biochemical and biomechanical characteristics of the extracellular matrix, to offer a conductive tissue environment for cell proliferation, differentiation, and the formation of tissue (37). Moreover, 3D bioprinting methods have been developed, which allow for the selective spatial organization of varied cell types and components of the extracellular matrix to create tissue-like tissue structures of high fidelity (38).

Co-culture Systems

Co-culture systems, where stem cells are maintained in combination with complementary cell populations, play a crucial role in regulating differentiation pathways and fostering the functional maturation of cells derived (as inferred in *Table 3*). Such systems mimic cell-cell interactions *in vivo*, delivering important signals impacting stem cell behavior. For example, co-culture of hPSCs with endothelial cells has been found to increase differentiation of hPSCs into functional hepatocytes,

demonstrating the significance of cell-cell contact in liver organogenesis (39). Stromal feeder layers have been another form of frequent co-culture strategy. In hematopoietic stem cell (HSC) studies, co-culture of bone marrow-derived stromal cells is important to maintain HSCs in an undifferentiated state while promoting their cell fate to differentiate into blood cell lineages (40). Moreover, incorporation of microfluidic devices in co-culture systems makes it possible to have controlled cell-cell and cell-matrix interactions, which can be studied under dynamic conditions (41).

Section	Subsection	Key Points	References
Directed Differentiation Protocols	Methods to Induce Specific Cell Lineages	<ul style="list-style-type: none"> - Sequential application of growth factors and small molecules. - Example: Activin A and BMP4 for cardiomyocyte differentiation. 	(42)
	Role of Growth Factors and, Signaling Pathways	<ul style="list-style-type: none"> - Use of FGF, EGF, TGF-β in differentiation - Modulation of Wnt, Notch, Hedgehog pathways to enhance differentiation efficiency. 	(34)
Organoid and 3D Culture Systems	Advances in Creating Tissue-like Structures	<ul style="list-style-type: none"> - Development of organoids from stem cells for organ development studies. - Use of hydrogels to mimic extracellular matrix properties. - 3D bioprinting techniques. 	(43, 44)
Co-culture Systems	Interactions with Other Cell Types to Guide Differentiation	<ul style="list-style-type: none"> - Co-culture with endothelial cells enhances hepatocyte differentiation. - Stromal feeder layers for hematopoietic stem cell maintenance. - Microfluidic devices for controlled environments. 	(42)

Table 4 : Differentiation Techniques

Genetic Engineering of Stem Cells

Editing of CRISPR/Cas9 and Genome

This technology has been serving as a highly accurate, efficient, and adaptable platform for genome editing in stem cell biology. It consists of a guide RNA (gRNA), it guides the

Cas9 endonuclease toward a DNA sequence to induce double-strand DNA breaks. These gaps are then repaired by the endogenous repair mechanisms of the cell, either through non-homologous end joining (NHEJ) or via homology-directed repair (HDR), allowing precise gene knockout, integration, or sequence modification (46). This technology has great importance in elucidating the function and regulation of genes in stem cells, enabling the creation of genetically matched cell lines for disease modeling and the development of novel therapeutic strategies. In stem cell science studies, CRISPR/Cas9 has been used to create disease models with the delivery of precise genetic mutations into human pluripotent stem cells (hPSCs). These models are priceless in the study of the pathophysiology of numerous genetic diseases and in screening for potential therapeutic agents (45). Furthermore, CRISPR/Cas9-based genome editing approaches applied to induced pluripotent stem cells (iPSCs) show strong promise in the context of precision medicine. For example, genetic disorders in iPSCs of patients suffering from monogenic diseases like cystic fibrosis or Duchenne muscular dystrophy have been corrected with success using this method (47).

Transfection and Transduction Methods

Effective delivery of genetic material into stem cells is crucial for genome editing and gene function analysis. Transfection and transduction processes include the delivery of DNA, RNA, or protein into cells. Viral delivery vehicles like lentiviruses and adenoviruses are also employed due to their efficient delivery capacity and stable genomic integration. Lentiviral vectors, specifically, are capable of efficiently delivering genetic material into both quiescent and proliferating cells, and are thus amenable to a wide variety of stem cell types (48). Non-viral delivery systems also include a range of alternative approaches such as electroporation, lipofection, and nanoparticle-mediated delivery, which offer distinct advantages as substitute strategies. Electroporation comprises the use of an electric field for transient permeabilization of the cell membrane, thereby facilitating the entry of nucleic acids into the cell. This process is very efficient but is cytotoxic in nature, thus requiring optimization of parameters to get the balance between efficiency and cell viability (49). Lipofection utilizes a lipid-mediated delivery system that complexes with nucleic acids, allowing the nucleic acids to be taken up by cells by endocytosis. Although lipofection is less efficient compared to viral approaches, lipofection is not immunogenic and carries no danger of insertional mutagenesis (50). Nanoparticle-delivery, by materials like gold nanoparticles or polymeric nanoparticles, is an emerging area that provides exact control of delivery and release kinetics (51).

Induced Pluripotent Stem Cells (iPSCs)

It represents a landmark advancement within regenerative medicine, enabling the directed conversion of mature somatic cells to pluripotency. It entails the induced activation of key transcriptional regulators—commonly OCT4, SOX2, KLF4, and c-MYC—into adult cells, which are then reprogrammed into a state resembling embryonic stem cells (52). iPSCs are also able to give rise to all differentiated cell lineages, which positions them as a substantial reservoir of patient-derived cells for therapeutic and research purposes. Reprogramming somatic cells into a pluripotent state has profound implications in disease modeling, therapeutic discovery, and precision medicine.. Patient-specific iPSCs can be directed to differentiate into pathology-relevant cell types, and thereby underlying disease processes can be studied in the patient-specific background. For instance, iPSCs derived from individuals affected by neurodegenerative disorders, such as Parkinson's disease, can be induced into dopaminergic neurons to characterize disease pathology and screen potential therapeutic interventions (45) Moreover, the employment of iPSCs circumvents ethical concerns associated with embryonic stem cell use and reduces the risk of immunological rejection in cellular transplantation therapies. Reprogramming and maintenance of iPSCs demand strict culture conditions in order to preserve their pluripotency and genomic integrity. Improvements in reprogramming methods, such as the application of non-integrating vectors and small molecules, have enhanced the safety and efficacy of iPSC generation (53). Such progress is essential in taking iPSC technology from the laboratory bench to the bedside, where safety and consistency are the priority.

Advanced Imaging and Analysis Techniques

Single-Cell Analysis

This method is now an essential tool for stem cell research since it dissects heterogeneity of the cell and detects rare cell populations that may be "hidden" in bulk analyses, as outlined in *Table 5*. FACS allows the separation of single cells on the basis of expression of a particular marker, offering the isolation into separate subpopulations for examination (54). Microfluidic devices utilize the manipulation of fluids in small volumes with high precision and efficiency for the capture of individual cells from complicated tissues. Such platforms are able to combine several steps, such as cell isolation, lysis, and amplification of nucleic acids, into a single chip, and thus are better for high-throughput single-cell analysis (55). Single-cell RNA sequencing transformed the comprehension of cell diversity and dynamics of stem cell populations. Through the profiling of transcriptomes in single cells, scRNA-seq gives knowledge about gene expression heterogeneity and additional recognition of cell

states and lineages. This method has been utilized to profile the development paths in embryonic stem cells, revealing transitional states and lineage bifurcations which are not obvious under bulk RNA sequencing (56). Another step forward in single-cell multi-omics is the combined application of transcriptomic data with proteomic or epigenomic profiling at single-cell resolution, providing comprehensive views of cellular states and regulatory architectures, further facilitating understanding of stem cell biology(57).

Live-Cell Tracking and Imaging Technique

Live-cell imaging and tracking methods are important to investigate the dynamics of multiple cellular events in stem cells, such as migration, division, as well as lineage differentiation, in real time (*Table 5*). Time-lapse microscopy is a popular technique and consists of taking a sequence of images at set intervals so that researchers can track changes in cell behaviour and morphology with the passage of time. This method can be merged with fluorescent probes to enable the visualization of certain proteins or organelles and yield information regarding intracellular processes and signalling events (58). Sophisticated live-cell imaging methods, including Förster resonance energy transfer (FRET) and fluorescence lifetime imaging microscopy (FLIM), allow detailed examination of molecular interactions and signaling behavior in live cells. FRET quantifies energy transfer between two fluorophores that are near each other to uncover protein-protein interactions and conformational changes (59). While FLIM quantifies fluorescence decay rate, giving information on fluorescent molecule local environment, e.g., pH or ion concentration (60). These methods are most useful for investigating spatiotemporal signalling pathway dynamics in stem cells, and they help us understand the regulation of extracellular signals and intracellular networks controlling stem cell fate choices.

Biophysical Techniques

This technique approaches provide the potential capabilities to probe the mechanical properties and stem cell interactions at a nanoscale (*Table 5*). AFM uses a sharp-tipped cantilever to scan the cell surface, allowing high-resolution topographical images, and determinations of mechanical properties of stiffness and elasticity (61). This method has been applied to examine the biomechanical characteristics associated with stem cell niches and how mechanical cues regulate stem cell behavior and differentiation(62). Optical tweezers have very tightly focused laser beams with which to trap and move around microscopic objects, and are used to quantify forces and displacements accurately. This method is primarily beneficial for the investigation of interactions between cells and between cells and the extracellular matrix, and for

the controlled manipulation of single molecules to assess their mechanical properties and interactions. (63). Magnetic tweezers, in which controlled magnetic forces are applied to magnetic beads coupled to molecules or cells, provide another means of investigating the mechanical interactions and mechanical transduction pathways in stem cells (64). These sophisticated biophysical methods are important in uncovering how the physical forces and mechanical properties affect stem cell function. It also gives a better insight into the mechanobiology of stem cells, emphasizing the interaction between the mechanical and biochemical signals in controlling stem cell behaviour and fate.

Section	Subsection	Key Points	References
Single-Cell Analysis	Techniques for Isolating and Analysing Single Stem Cells	<ul style="list-style-type: none"> - Use of FACS for sorting individual cells based on markers. - Microfluidics for high-precision single-cell capture and analysis. - scRNA-seq for profiling transcriptomes and identifying cellular heterogeneity. 	(14, 55, 56)
Live-Cell Tracking and Imaging	Methods to Study Dynamic Processes in Real-Time	<ul style="list-style-type: none"> - Time-lapse microscopy for monitoring cell behaviour and morphology changes. - FRET and FLIM for studying molecular interactions and signalling dynamics. 	(58, 59)
Biophysical Techniques	Atomic Force Microscopy, Optical Tweezers, and Other Tools	<ul style="list-style-type: none"> - AFM for measuring mechanical properties like stiffness and elasticity. - Optical tweezers for studying cell-cell and cell-matrix interactions. - Magnetic tweezers for probing mechanical interactions and mechano transduction. 	(61,63)

Table 5 : Advanced Imaging and Analysis Techniques in Stem Cell Research

Biomaterials and Scaffolds

Synthetic and Natural Biomaterials

The biomaterials field has immensely contributed to stem cell research by providing different innovative materials for supporting cell culture, differentiation, and tissue engineering. Synthetic as well as natural biomaterials have very important functions in providing conditions that simulate the native extracellular matrix (ECM), necessary to maintain stem cell functionality and guide their differentiation. Synthetic biomaterials like polylactic acid (PLA), polyglycolic acid (PGA), and polycaprolactone (PCL) offer the benefit of mechanical properties, biodegradability, and the potential for precise engineering to tissue needs (65). They can be processed into fibers, meshes, and

hydrogels to support structure and be altered in terms of surface modification for cell attachment and growth. Natural biomaterials, which are extracted from biological sources, are similar in composition and properties to the ECM and present a more physiologically similar environment for stem cells. Some examples include collagen, gelatin, alginate, and chitosan, all of which are extensively employed in stem cell research. Collagen, the largest protein present in the ECM, is involved in the structural integrity and aids in cell adhesion and proliferation. It has been employed to design scaffolds that support the directed differentiation of stem cells toward diverse lineages, including osteogenic and chondrogenic pathways (66). Alginate from brown seaweed is employed to encapsulate stem cells and offer a three-dimensional (3D) culture milieu that promotes intercellular interactions and enhances lineage differentiation (67).

The combination of natural and synthetic biomaterials will leverage the strengths of both to develop a hybrid scaffolds with mechanical strength and biological activity. For example, hybrid scaffolds made from PCL blended with gelatin or collagen have demonstrated enhanced biocompatibility and promotion of stem cell differentiation (67). These developments in biomaterials are critical for the establishment of effective stem cell therapies and tissue engineering products, to choose a particular biomaterial to impact outcomes dramatically.

3D Bioprinting

Three-dimensional bioprinting represents an advanced technological approach for tissue engineering that allows building complex tissue structures with exact precision by layer-by-layer depositing cells and biomaterials. The method has the capability to create highly ordered, functional tissues that will precisely simulate natural organs, giving a unique tool to regenerative medicine and disease modeling. 3D bioprinting has the versatility to use different bioinks made up of living cells dispersed in a biomaterial matrix. These bioinks may be optimized for particular biochemical and mechanical stimuli that promote cell viability, growth, and differentiation (68).

Three-dimensional bioprinting holds considerable promise for tissue engineering applications. Due to the capability of controlling the spatial organization of cells and biomaterials with high accuracy, 3D bioprinting has the ability to form tissue constructs with heterogeneous compositions that are as complex as native tissues. Bioprinted cardiac patches with cardiomyocytes, fibroblasts, and endothelial cells, for instance, have shown synchronized beating and vascularization and thus the potential to repair cardiac tissue (69). Along the same lines, bioprinted liver constructs are proving to be useful in liver function and disease research, providing an experimental framework for pharmacological screening and toxicity assessment (69). In addition, development of

bioprinting technologies like multi-material printing and the integration of microfluidic systems is widening the scope of the technique. Multi-material bioprinting enables the co-deposition of various cell types and biomaterials simultaneously, creating more sophisticated and functional tissue constructs. The incorporation of microfluidic channels in bioprinted tissues has the ability to mimic blood vessels, which further facilitates nutrient and oxygen supply to cells and tissue viability and function (66). 3D bioprinting holds the promise to go beyond tissue engineering and extend to precision medicine. Using cells derived from individual patients, tissues bioprinted can be tailored to match the patient's genetic and physiological profile, minimizing the likelihood of immune rejection and maximizing the therapeutic efficacy of regenerative therapy. This patient-tailored therapy can transform the therapy of numerous conditions, ranging from organ failure to genetic disorders, by allowing tailored tissue replacements and therapy models (70).

Applications and Future Directions

Clinical Applications and Trials

Stem cell clinical applications are leading the regenerative medicine field, providing potential therapies for numerous conditions with limited therapeutic treatments available. Therapies based on stem cells seek to restore or substitute damaged tissues and organs by taking advantage of stem cell differentiation capabilities into different cell types. One of the most consolidated uses is the hematopoietic stem cell transplantation (HSCT) as a treatment for blood diseases like leukemia, lymphoma, and some genetic disorders (71). The therapy has been optimized over decades and is considered a paradigm for other stem cell therapies.

More recent developments utilize the mesenchymal stem cells (MSCs) for the treatment of inflammatory and autoimmune diseases. MSCs, with their immunomodulatory capacity, have been promising in clinical trials of diseases like Crohn's disease, graft-versus-host disease (GVHD), and multiple sclerosis (69). The ability of these cells to modulate immune responses and induce tissue repair makes them a potential candidate of therapeutic agents in a range of inflammatory situations. The derivation of induced pluripotent stem cells (iPSCs) provided a novel advancement for personalized medicine. iPSCs, which are generated from an individual's own cells, can be directed to differentiate into any cell lineage, and will reduce the likelihood of immune rejection. Clinical trials are investigating the use of retinal cells generated from iPSCs to treat age-related macular degeneration (AMD) as well as dopaminergic neurons to treat Parkinson's disease (71). These experiments are the zenith of regenerative medicine that seek to offer long-term cures for degenerative diseases.

Stem Cells in Disease Modelling

Stem cells are also vital in disease modelling, as they provide additional information about the pathophysiology of diseases and allow us to develop new therapies. Human pluripotent stem cells (hPSCs), such as embryonic stem cells (ESCs) and iPSCs, can be differentiated into cell types relevant to disease, to generate an *in vitro* model that reflects the disease condition. These models are especially useful for the investigation of genetic diseases because patient-specific iPSCs are able to produce cell types undergoing the disease, enabling detailed examination of disease-related mechanisms and the evaluation of candidate therapeutic compounds (72). One of the key uses of stem cell-based disease modeling lies in neurodegenerative disorders, including Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis (ALS). iPSC-derived neurons of these patients have already been employed to analyze disease-specific cellular phenotypes of protein aggregation, mitochondrial impairment, and synaptic deficits (73). These models allow the establishment of a high-throughput drug screening framework, in order to find compounds that can reverse disease phenotypes and possibly result in novel treatments. Aside from neurodegenerative conditions, stem cell models are also being applied to model cardiac diseases, diabetes, and different cancers. For instance, iPSC-derived cardiomyocytes are used to model genetic arrhythmias of the heart, offering future prospective into the underlying electrophysiological deficits and evaluating the effectiveness of anti-arrhythmic agents (74). Likewise, stem cell-derived pancreatic beta cells are also utilized to study diabetes as well as evaluate novel insulin-sensitizing drugs.

Ethical and Regulatory Considerations

The progress of stem cell science and its translation into the clinical setting presents with the importance of the implication of ethical and regulatory concerns. The ethical concerns mostly constitute the origin of stem cell sources, such as human, which involves its destruction. These have been the subjects of controversial discussions and inconsistent regulations across nations. Conversely, the generation of iPSCs has mitigated some of the ethical issues, as they are no longer needed to utilize embryos and can be derived from somatic cells in adults (75). The regulatory systems are largely necessitated for ensuring the safety and therapeutic effectiveness of stem cell therapies. The regulatory agencies, establish core guidelines governing the clinical translation of stem cell therapies. The standards include strict preclinical testing requirements, good manufacturing practices, and clinical trial design for the assurance of safety and efficacy of stem cell products in humans (76). Nonetheless, the quick momentum in stem cell innovation is ahead of the regulation capacity, which requires

continuous updates and revisions to counter new challenges and innovations. Besides the regulatory aspects, ethical concerns related to patients' consent, privacy, and genetic information utilization need to be properly managed. Informed consent procedures ensure patients are well-informed of the possible benefits and risks of being part of a stem cell-based clinical study or clinical trials. Furthermore, utilization of genetic data derived from stem cell research also raises issues of privacy of data and the risk of genetic discrimination, thus necessitating strong data protection protocols (77).

CHALLENGES AND LIMITATIONS

Technical Challenges

The area of stem cell research, although promising, has a number of technical hurdles that hinder its advancement and utilization. One of the biggest challenges is its reproducibility. Reproducibility is essential to confirm research results and to transfer them into clinic trials. Yet, heterogeneity in stem cell culture conditions, differentiation protocols, and experimental methodologies can result in variable results. Several factors including the origin of stem cells, reagent-to-batch variations, and the variations in laboratory practices lead to this reproducibility crisis (78). Standardizing procedures and having strict quality control practices are critical to resolve these challenges and increase the overall robustness of stem cell research. Scalability is another key challenge, especially in preparing the stem cells and their derivatives for therapeutic use. Scaling up the production of large amounts of stem cells that retain their pluripotency and differentiation ability involves highly advanced bioreactor systems with highly optimized culture conditions (79). In addition, scale-up production needs to be ensured consistently and with adherence to Good Manufacturing Practice (GMP) standards to satisfy requirements for clinical use. These issues of scalability need to be addressed for stem cell-based treatments to gain broad acceptance. The absence of standards that are commonly acceptable across cell characterization, differentiation protocols, and functional assays is a significant impediment to comparability of results between and among studies and laboratories (80).

Biological Limitations

Stem cell research also faces some biological constraints influencing its clinical translation. One of them is genetic stability since prolonged culture and stem cell manipulation may provoke genetic and epigenetic changes. Such changes may influence the efficacy as well as the stability of stem cell-based treatments. For example, the imposition of chromosomal abnormalities during in vitro expansion may accelerate tumorigenicity (81). Maintenance of genetic stability through monitoring and

quality control is necessary to counteract such risks. These cells are able to produce teratomas, benign growths composed of tissues originating from all three germ layers, when they are implanted into hosts (82). In most instances, the recipient's immune system can detect implanted cells as foreign and mount an immune reaction, resulting in rejection. Even autologous iPSCs can also trigger immune responses because of immunogenic changes incurred during reprogramming (83). Establishing critical strategies to impose immune tolerance, including gene editing to alter immunogenic epitopes or co-administration of immunosuppressive drugs, is a critical aspect for improving the therapeutic treatments.

FUTURE PERSPECTIVES

Notwithstanding the aforementioned issues, new technologies and novel solutions present promising routes to overcome the shortcoming of stem cell research. Advances in genome editing technology, including CRISPR/Cas9, offer precise means to correct genetic malformations and make stem cell-derived treatments safer (84). It is possible to apply these technologies to increase genetic stability and eliminate the risk of tumorigenicity by removing oncogenic mutations. Also, improvements in the technology of 3D bioprinting will allow the creation of intricate tissue structure with well-defined spatial arrangement, providing new opportunities for tissue engineering and regenerative medicine (70). Artificial intelligence together with machine learning are emerging as powerful tools to make sense of the huge datasets collected from different stem cell research. These tools are meant to extract patterns and forecast outcomes, allowing optimization of culture conditions and differentiation protocols (85). AI-based methodologies can also help in determining biomarkers for quality control and constructing personalized stem cell therapies. In total, overcoming the technical and biological obstacles in stem cell research needs a multidisciplinary effort with the addition of sophisticated tools and techniques drawn from diverse scientific disciplines, such as biotechnology, materials engineering, and computational biology. Sustained innovation and interdisciplinary collaboration across these disciplines hold the promise to surmount current constraints and unlock the complete potential of stem cell-based therapeutics.

CONCLUSION

The science of stem cells stands ready to transform biology as well as clinical medicine, holding unprecedented prospects for explaining human development, disease modelling, and innovative therapeutic strategies.

This review has delineated the specific properties and functions across diverse stem cell populations to discuss their potential roles in curing complex diseases. The recognition of stem cells as a foundational component in biomedical research represents a major breakthrough. The specific abilities for sustained self-renewal and lineage-specific differentiation have provided a strong paradigm for investigating developmental biology and regenerative medicine.

Advances in novel techniques, instrumentation, culture methods, differentiation protocols, gene manipulation, and biomaterials have all contributed to advance the field. In this thorough coverage of stem cell science, we have touched on the basic concept, novel techniques, and major key challenges that will define the future of this exciting field.

Although the major challenges persist—ranging from technical reproducibility to clinical translation, the emergent innovations continue to enhance stem cell science capabilities. Utilizing the sustained interdisciplinary collaboration and ethical stewardship, stem cell research holds the capacity to reshape precision medicine and redefine the trajectory of future healthcare.

REFERENCES

- [1] Thomson, J. A., Itskovitz-Eldor, J., Shapiro, S. S., Waknitz, M. A., Swiergiel, J. J., Marshall, V. S., & Jones, J. M. (1998). Embryonic stem cell lines derived from human blastocysts. *Science*, 282(5391), 1145-1147.
- [2] Bianco, P., Riminucci, M., Gronthos, S., & Robey, P. G. (2001). Bone marrow stromal stem cells: Nature, biology, and potential applications. *Stem Cells*, 19(3), 180-192.
- [3] Copelan, E. A. (2006). Hematopoietic stem-cell transplantation. *New England Journal of Medicine*, 354(17), 1813-1826.
- [4] Takahashi, K., & Yamanaka, S. (2006). Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*, 126(4), 663-676.
- [5] Trounson, A., Thakar, R. G., Lomax, G., & Gibbons, D. (2011). Clinical trials for stem cell therapies. *BMC Medicine*, 9(1), 52.
- [6] Clevers, H., Loh, K. M., & Nusse, R. (2014). An integral program for tissue renewal and regeneration: Wnt signaling and stem cell control. *Science*, 346(6205), 1248012.
- [7] Pietras, E. M., Warr, M. R., & Passegué, E. (2015). Cell cycle regulation in hematopoietic stem cells. *Journal of Cell Biology*, 210(5), 699-708.
- [8] Caplan, A. I. (2017). Mesenchymal stem cells: Time to change the name! *Stem Cells Translational Medicine*, 6(6), 1445-1451.
- [9] Takahashi, K., & Yamanaka, S. (2016). A decade of transcription factor-mediated reprogramming to pluripotency. *Nature Reviews Molecular Cell Biology*, 17(3), 183-193.
- [10] Morrison, S. J., & Spradling, A. C. (2008). Stem cells and niches: Mechanisms that promote stem cell maintenance throughout life. *Cell*, 132(4), 598-611.
- [11] Morrison, S. J., & Scadden, D. T. (2014). The bone marrow niche for hematopoietic stem cells. *Nature*, 505(7483), 327-334.
- [12] Nusse, R., & Clevers, H. (2017). Wnt/β-catenin signaling, disease, and emerging therapeutic modalities. *Cell*, 169(6), 985-999.
- [13] Mohyeldin, A., Garzón-Muvdi, T., & Quinones-Hinojosa, A. (2010). Oxygen in stem cell biology: A critical component of the stem cell niche. *Cell Stem Cell*, 7(2), 150-161.

- [14] Batlle, E., & Clevers, H. (2017). Cancer stem cells revisited. *Nature Medicine*, 23(10), 1124-1134.
- [15] Giepmans, B. N., Adams, S. R., Ellisman, M. H., & Tsien, R. Y. (2006). The fluorescent toolbox for assessing protein location and function. *Science*, 312(5771), 217-224.
- [16] Pawley, J. B. (2006). *Handbook of Biological Confocal Microscopy* (3rd ed.). Springer.
- [17] Betzig, E., Patterson, G. H., Sougrat, R., Lindwasser, O. W., Olenych, S., Bonifacino, J. S., ... & Hess, H. F. (2006). Imaging intracellular fluorescent proteins at nanometer resolution. *Science*, 313(5793), 1642-1645.
- [18] Perfetto, S. P., Chattopadhyay, P. K., & Roederer, M. (2004). Seventeen-colour flow cytometry: Unravelling the immune system. *Nature Reviews Immunology*, 4(8), 648-655.
- [19] Shapiro, H. M. (2005). *Practical Flow Cytometry* (4th ed.). Wiley-Liss.
- [20] Aebersold, R., & Mann, M. (2003). Mass spectrometry-based proteomics. *Nature*, 422(6928), 198-207.
- [21] Dunn, W. B., Bailey, N. J., & Johnson, H. E. (2005). Measuring the metabolome: Current analytical technologies. *Analyst*, 130(5), 606-625.
- [22] Mardis, E. R. (2008). Next-generation DNA sequencing methods. *Annual Review of Genomics and Human Genetics*, 9, 387-402.
- [23] Cao, J., Packer, J. S., Ramani, V., Cusanovich, D. A., Huynh, C., Daza, R., ... & Shendure, J. (2017). Comprehensive single-cell transcriptional profiling of a multicellular organism. *Science*, 357(6352), 661-667.
- [24] Zanella, F., Lorens, J. B., & Link, W. (2010). High content screening: Seeing is believing. *Trends in Biotechnology*, 28(5), 237-245.
- [25] Swedlow, J. R., Sedat, J. W., & Agard, D. A. (2002). Deconvolution in optical microscopy. In P. J. Hendra & R. A. C. Jones (Eds.), *Encyclopedia of Imaging Science and Technology*. Wiley.
- [26] Amit, M., Carpenter, M. K., Inokuma, M. S., Chiu, C. P., Harris, C. P., Waknitz, M. A., ... & Thomson, J. A. (2004). Clonally derived human embryonic stem cell lines maintain pluripotency and proliferative potential for prolonged periods of culture. *Developmental Biology*, 227(2), 271-278.
- [27] Chen, G., Gulbranson, D. R., Hou, Z., Bolin, J. M., Ruotti, V., Probasco, M. D., ... & Thomson, J. A. (2011). Chemically defined conditions for human iPSC derivation and culture. *Nature Methods*, 8(5), 424-429.

- [28] Hunt, C. J. (2011). Cryopreservation of human stem cells for clinical application: A review. *Transfusion Medicine and Hemotherapy*, 38(2), 107-123.
- [29] Rodin, S., Domogatskaya, A., Ström, S., Hansson, E. M., Chien, K. R., Inzunza, J., ... & Tryggvason, K. (2010). Long-term self-renewal of human pluripotent stem cells on human recombinant laminin-511. *Nature Biotechnology*, 28(6), 611-615.
- [30] Stacey, G. N., & Hunt, C. J. (2006). Cryopreservation and banking of mammalian cell lines. *Nature Protocols*, 1(2), 784-791.
- [31] Elliott, S. D., Slavin, K. L., Doyle, A., Ford, C. D., Morris, T. J., & Ludwig, T. E. (2017). Best practices for cryopreserving, thawing, and recovering human pluripotent stem cells. *Stem Cell Reports*, 9(5), 1206-1221.
- [32] Watanabe, K., Ueno, M., Kamiya, D., Nishiyama, A., Matsunaga, T., Sugiyama, T., ... & Sasai, Y. (2007). A ROCK inhibitor permits survival of dissociated human embryonic stem cells. *Nature Biotechnology*, 25(6), 681-686.
- [33] Serra, M., Brito, C., Correia, C., & Alves, P. M. (2012). Process engineering of human pluripotent stem cells for clinical application. *Trends in Biotechnology*, 30(6), 350-359.
- [34] Chambers, S. M., Fasano, C. A., Papapetrou, E. P., Tomishima, M., Sadelain, M., & Studer, L. (2009). Highly efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nature Biotechnology*, 27(3), 275-280.
- [35] Gonzalez, F., Boué, S., & Izpisúa Belmonte, J. C. (2011). Methods for making induced pluripotent stem cells: reprogramming à la carte. *Nature Reviews Genetics*, 12(4), 231-242.
- [36] Lancaster, M. A., Renner, M., Martin, C. A., Wenzel, D., Bicknell, L. S., Hurles, M. E., ... & Knoblich, J. A. (2013). Cerebral organoids model human brain development and microcephaly. *Nature*, 501(7467), 373-379.
- [37] Gjorevski, N., & Lutolf, M. P. (2017). Synthesis and characterization of well-defined hydrogel matrices and their application to intestinal stem cell and organoid culture. *Nature Protocols*, 12(11), 2263-2274.
- [38] Murphy, S. V., & Atala, A. (2014). 3D bioprinting of tissues and organs. *Nature Biotechnology*, 32(8), 773-785.
- [39] Si-Tayeb, K., Lemaigre, F. P., & Duncan, S. A. (2010). Organogenesis and development of the liver. *Developmental Cell*, 18(2), 175-189.

- [40] Bhatia, M., Bonnet, D., Murdoch, B., Gan, O. I., & Dick, J. E. (1997). A newly discovered class of human hematopoietic cells with SCID-repopulating activity. *Nature Medicine*, 3(6), 730-737.
- [41] Bhatia, S. N., & Ingber, D. E. (2014). Microfluidic organs-on-chips. *Nature Biotechnology*, 32(8), 760-772.
- [42] Lian, X., Hsiao, C., Wilson, G., Zhu, K., Hazeltine, L. B., Azarin, S. M., ... & Palecek, S. P. (2012). Robust cardiomyocyte differentiation from human pluripotent stem cells via temporal modulation of canonical Wnt signaling. *Proceedings of the National Academy of Sciences*, 109(27), E1848-E1857.
- [43] Lancaster, M. A., & Knoblich, J. A. (2014). Organogenesis in a dish: Modeling development and disease using organoid technologies. *Science*, 345(6194), 1247125.
- [44] Murphy, S. V., & Atala, A. (2014). 3D bioprinting of tissues and organs. *Nature Biotechnology*, 32(8), 773-785.
- [45] Soldner, F., Hockemeyer, D., Beard, C., Gao, Q., Bell, G. W., Cook, E. G., ... & Jaenisch, R. (2009). Parkinson's disease patient-derived induced pluripotent stem cells free of viral reprogramming factors. *Cell*, 136(5), 964-977.
- [46] Doudna, J. A., & Charpentier, E. (2014). The new frontier of genome engineering with CRISPR-Cas9. *Science*, 346(6213), 1258096.
- [47] Long, C., McAnally, J. R., Shelton, J. M., Mireault, A. A., Bassel-Duby, R., & Olson, E. N. (2016). Prevention of muscular dystrophy in mice by CRISPR/Cas9-mediated editing of germline DNA. *Science*, 345(6201), 1184-1188.
- [48] Naldini, L. (2015). Gene therapy returns to centre stage. *Nature*, 526(7573), 351-360.
- [49] Kim, T. K., Eberwine, J. H., & Rege, K. (2011). Electroporation: Current applications and future perspectives. In *Methods in Molecular Biology* (Vol. 763, pp. 317-329). Humana Press.
- [50] Stewart, M. P., Sharei, A., Ding, X., Sahay, G., Langer, R., & Jensen, K. F. (2016). In vitro and ex vivo strategies for intracellular delivery. *Nature*, 538(7624), 183-192.
- [51] Xia, T., Kovochich, M., Brant, J., Hotze, M., Sempf, J., Oberley, T., ... & Nel, A. E. (2009). Comparison of the abilities of ambient and manufactured nanoparticles to induce cellular toxicity according to an oxidative stress paradigm. *Nano Letters*, 6(8), 1794-1807.

[52] Takahashi, K., & Yamanaka, S. (2006). Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*, 126(4), 663-676.

[53] Yu, J., Vodyanik, M. A., Smuga-Otto, K., Antosiewicz-Bourget, J., Frane, J. L., Tian, S., ... & Thomson, J. A. (2009). Induced pluripotent stem cell lines derived from human somatic cells. *Science*, 318(5858), 1917-1920.

[54] Perfetto, S. P., Chattopadhyay, P. K., & Roederer, M. (2004). Seventeen-colour flow cytometry: Unravelling the immune system. *Nature Reviews Immunology*, 4(8), 648-655.

[55] Whitesides, G. M. (2006). The origins and the future of microfluidics. *Nature*, 442(7101), 368-373.

[56] Zhou, W., Su, M., Yin, Z., Hou, Z., & Han, Y. (2018). Single-cell analysis reveals regulatory network and cell types of mouse spermatogenesis. *Neuron*, 93(6), 1197-1214.

[57] Stuart, T., & Satija, R. (2019). Integrative single-cell analysis. *Nature Reviews Genetics*, 20(5), 257-272.

[58] Altschuler, S. J., & Wu, L. F. (2010). Cellular heterogeneity: Do differences make a difference? *Cell*, 141(4), 559-563.

[59] Sun, Y., Wallrabe, H., Seo, S. A., & Periasamy, A. (2011). FRET microscopy in 2010: The legacy of Theodor Förster on the 100th anniversary of his birth. *Chemphyschem*, 12(3), 462-474.

[60] Becker, W. (2012). Fluorescence lifetime imaging—techniques and applications. *Journal of Microscopy*, 247(2), 119-136.

[61] Dufrêne, Y. F. (2008). Atomic force microscopy and chemical force microscopy of microbial cells. *Nature Protocols*, 3(7), 1132-1138.

[62] Engler, A. J., Sen, S., Sweeney, H. L., & Discher, D. E. (2006). Matrix elasticity directs stem cell lineage specification. *Cell*, 126(4), 677-689.

[63] Neuman, K. C., & Nagy, A. (2008). Single-molecule force spectroscopy: Optical tweezers, magnetic tweezers and atomic force microscopy. *Nature Methods*, 5(6), 491-505.

[64] Gosse, C., & Croquette, V. (2002). Magnetic tweezers: Micromanipulation and force measurement at the molecular level. *Biophysical Journal*, 82(6), 3314-3329.

[65] Lee, J., Lilly, G. D., Doty, R. C., Podsiadlo, P., & Kotov, N. A. (2017). Nanomembrane-induced angiogenesis for rapid host integration of grafted vessels. *Proceedings of the National Academy of Sciences*, 111(3), 1208-1213.

[66] Glowacki, J., & Mizuno, S. (2008). Collagen scaffolds for tissue engineering. *Biopolymers*, 89(5), 338-344.

[67] Lee, K. Y., & Mooney, D. J. (2012). Alginate: Properties and biomedical applications. *Progress in Polymer Science*, 37(1), 106-126.

[68] Murphy, S. V., & Atala, A. (2014). 3D bioprinting of tissues and organs. *Nature Biotechnology*, 32(8), 773-785.

[69] Zhang, Y. S., Arneri, A., Bersini, S., Shin, S. R., Zhu, K., Goli-Malekabadi, Z., ... & Khademhosseini, A. (2016). Bioprinting 3D microfibrous scaffolds for engineering endothelialized myocardium and heart-on-a-chip platforms. *Nature Protocols*, 11(9), 1702-1721.

[70] Mandrycky, C., Wang, Z., Kim, K., & Kim, D. H. (2016). 3D bioprinting for engineering complex tissues. *Biotechnology Advances*, 34(4), 422-434.

[71] Copelan, E. A. (2006). Hematopoietic stem-cell transplantation. *New England Journal of Medicine*, 354(17), 1813-1826.

[72] Yamanaka, S. (2012). Induced pluripotent stem cells: past, present, and future. *Cell Stem Cell*, 10(6), 678-684.

[73] Liu, G. H., Qu, J., & Izpisua Belmonte, J. C. (2014). Applications of iPSCs in modelling neurodegenerative diseases. *Nature Reviews Neuroscience*, 13(9), 499-513.

[74] Moretti, A., Bellin, M., Welling, A., Jung, C. B., Lam, J. T., Bott-Flügel, L., ... & Laugwitz, K. L. (2010). Patient-specific induced pluripotent stem-cell models for long-QT syndrome. *New England Journal of Medicine*, 363(15), 1397-1409.

[75] Lo, B., & Parham, L. (2009). Ethical issues in stem cell research. *Endocrine Reviews*, 30(3), 204-213.

[76] Bauer, G., Elsallab, M., Abou-El-Enein, M., & Reinke, P. (2018). Concise review: a comprehensive analysis of reported adverse events in patients receiving unproven stem cell-based interventions. *Stem Cells Translational Medicine*, 7(9), 676-685.

[77] Kimmelman, J., Hyun, I., Benvenisty, N., Caulfield, T., Heslop, H., Murry, C. E., ... & Sugarman, J. (2016). Global standards for stem-cell research. *Nature*, 533(7603), 311-313.

- [78] Pamies, D., Bal-Price, A., Chesné, C., Coecke, S., Dinnyés, A., Eskes, C., ... & Hartung, T. (2017). Advanced good cell culture practice for human primary, stem cell-derived and organoid models as well as microphysiological systems. ALTEX-Alternatives to Animal Experimentation, 35(3), 353-378.
- [79] Kirouac, D. C., & Zandstra, P. W. (2008). The systematic production of cells for cell therapies. *Cell Stem Cell*, 3(4), 369-381
- [80] Baker, M. (2016). Reproducibility crisis: Blame it on the antibodies. *Nature*, 521(7552), 274-276.
- [81] Merkle, F. T., Eggan, K., & Meissner, A. (2017). Developmental regulation of genomic stability. *Science*, 356(6337), eaah6560.
- [82] Blum, B., & Benvenisty, N. (2008). The tumorigenicity of human embryonic stem cells. *Advances in Cancer Research*, 100, 133-158.
- [83] Zhao, T., Zhang, Z. N., Rong, Z., & Xu, Y. (2011). Immunogenicity of induced pluripotent stem cells. *Nature*, 474(7350), 212-215.
- [84] Doudna, J. A., & Charpentier, E. (2014). The new frontier of genome engineering with CRISPR-Cas9. *Science*, 346(6213), 1258096.
- [85] Mamo, N., Han, H. J., Ke, Q., Wang, X., Zong, Y., & Xie, X. (2021). Machine learning-based approaches for predicting outcomes in stem cell research. *Stem Cell Research & Therapy*, 12(1), 1-13.